

UNPARTICLES AS FIELDS WITH CONTINUOUSLY DISTRIBUTED MASSES

N.V.Krasnikov

INR, Moscow

OUTLINE

1. Introduction
2. Fields with continuously distributed masses
3. Phenomenological implications
4. Conclusion

1. Introduction

Recently H.Georgi proposed a model of unparticles. The main speculation : suppose there is conformal invariant world (gauge theory with fermions with ultraviolet fixed point as an example). For such conformal world all 2-point functions $\langle O(p)O(-p) \rangle$ behave like $D(p^2) \sim (p^2)^{2\beta+2d-3}$, where β is anomalous dimension of the operator O and d its naive dimension. As a consequence there is no single particle pole in the spectrum . The spectrum is continuous.

The main support of the possible existence of 4-dimensional conformal models is due to the fact that for some number of matter fields in gauge models one-loop beta function contribution is negative while two-loop correction is positive that leads to speculation about existence of fixed point

For instance for QCD with n_f flavours

$$\beta(\alpha_s) = -\beta_0 \alpha_s^2 (2\pi)^{-1} - \beta_1 \alpha_s^3 (2\pi)^{-3} + O(\alpha_s^5)$$

$$\beta_0 = 11 - 2n_f/3; \quad \beta_1 = 51 - 19n_f/3$$

For $8 < n_f < 16$ in PT we have fixed point.

Suppose conformal “unparticle world” and our world are connected due to nonrenormalizable interaction

$$L_i = c\Lambda^{-n}O(\text{particle})O(\text{unparticle})$$

Due to assumed interactions two types of observable effects are possible:

a. Production of unparticles at colliders :

$$qq \rightarrow qU$$

$$gg \rightarrow gU$$

Unparticles being weakly interacting in our world are not detected and behave like neutrino

As a consequence we obtain that unparticles signature are events with missing transverse momentum and hadronic jet(s) like in ADD model which describes our 4 dimensional world plus gravity in $(4 + n)$ dimensional world with compactification of n additional dimensions (tower of massive gravitons).

b. Exchange of unparticles leads to additional propagators

$$D(p^2) \sim (p^2)^{-\Delta}$$

that change the SM predictions for cross sections like $DY, \gamma\gamma$ production, ...

In other words:

- ***Due to assumed interactions of particles and unparticles it is possible to produce unparticles in particle collisions .***
- ***As a consequence of continuous spectrum of unparticles and weak interactions with particles unparticles are not detected . How to detect unparticles?***

1. Missing Transverse Energy

2. Unparticle exchange leads to the modification of particle propagators. So study of processes like dimuon production allows to constrain particle-unparticle interactions

Some unparticle references:

1.H.Georgi, Phys.Rev.Lett. 98 221601(1997).

Plus a lot of “unparticle exercises” , for instance:

2. K.Cheung et al, Collider Phenomenology of Unparticle Physics, arXiv:07063155

In this talk I show that the notion of an unparticle can be described as a particular case of a field with continuously distributed mass. I also review the models with continuously distributed masses and describe possible phenomenological implications for Large Hadron Collider(LHC)

This talk is based on my papers:

1. N.V.K., Higgs boson with continuously distributed mass, Phys.Lett. B325(1994)430.
2. N.V.K., Unparticle as a field with continuously distributed mass, Int.J.Mod.Phys. 22 (2007) 5117.
3. N.V.K., LHC signatures for Z' models with continuously distributed mass, Mod.Phys.Lett. 23 (2008) 3233.

Also I have to mention related work

A.A.Slavnov, Theor.Math.Phys. 148(2006)339

2. Fields with continuously distributed mass

Let us start with N free scalar fields $\Phi_k(x_k)$ with masses m_k . For the field

$$\Phi(x_k, m_k, c_k) = \sum c_j \Phi_j(x_j, m_j)$$

free propagator has the form

$$D_{\text{int}}(k^2) = \sum |c_j|^2 (k^2 - m_j^2 + i\varepsilon)^{-1} = \int \rho(t, c_j, m_j) (k^2 - t + i\varepsilon)^{-1} dt, \\ \rho(t, c_j, m_j) = \sum |c_j|^2 \delta(t - m_j^2),$$

n2

In the limit $N \rightarrow \infty$,

$$\rho(t, c_j, m_j) \rightarrow \rho(t) \text{ and} \\ D_{\text{int}}(k^2) \rightarrow \int \rho(t) (k^2 - t - i\varepsilon)^{-1} dt$$

For instance for

$$m_k^2 = m^2 + k\Delta N^{-1} \text{ and } |c_k|^2 = N^{-1} \text{ in the limit } N \rightarrow \infty \\ \rho(t) = \theta(t - m^2) \theta(m^2 + \Delta - t) \Delta^{-1}$$

For spectral density $\rho(t) \sim t^{\delta-1}$ the propagator

$$D_{\text{int}}(k^2) \sim (k^2)^{\delta-1}$$

that corresponds to the case of unparticle propagator and the limiting field $\Phi(x, \rho(t)) = \lim_{N \rightarrow \infty} \Phi(x, m_j, c_j)$ describes unparticle field. It is possible to introduce self interaction in standard way as

$$L_{\text{int}} = -\lambda(\Phi(x, \rho(t)))^4$$

For finite $\int \rho(t) dt$ the asymptotics of the effective propagator coincides with free propagator

$D(p) \sim (p^2)^{-1}$ and the model is renormalizable.

The generalization to the case of vector fields is straightforward. Consider the Stueckelberg Lagrangian

$$L = \sum_k [(-1/4) F_{\mu\nu, k} F^{\mu\nu, k} + (1/2) m_k^2 (A_{\mu, k} - \partial_\mu \Phi_k)^2]$$

Gauge invariance:

$$A_{\mu, k} \rightarrow A_{\mu, k} + \partial_\mu \alpha_k,$$

$$\Phi_k \rightarrow \Phi_k + \alpha_k$$

Слайд 10

п3

krasnikov; 29.06.2009

For the field $B_\mu = \sum c_k A_{\mu,k}$ in the limit

$N \rightarrow \infty$ we obtain free unparticle vector field for power like spectral density

One can introduce gauge invariant interaction with fermion field ψ in standard way

$$L_{\text{int}} = e \psi \gamma_\mu \psi B_\mu$$

For such model Feynman rules the same as in QED except the change photon propagator $1/k^2 \rightarrow D_{\text{int}}(k^2)$.

Another approach to the fields with continuously distributed mass related with

the introduction of additional space dimensions.

The main peculiarity is that we postulate Poincare invariance only in 4-dimensional space-time but not Poincare invariance in $(4+n)$ -dimensional space-time.

Consider scalar field $\Phi(x_\mu, x_4)$ in five-dimensional field interacting with the four-dimensional fermion field $\psi(x)$.

Слайд 11

n4

krasnikov; 29.06.2009

The scalar action has the form

$$S_1 = (1/2) \int [\partial_\mu \Phi \partial^\mu \Phi - \Phi f(-\partial_4^2) \Phi] d^5x$$

This action is invariant only under 4-dimensional Poincare group and 5-dimensional free propagator is

$$D_0 = (k_\mu k^\mu - f(k_4^2))^{-1}$$

The interaction of 5-dimensional scalar field with 4-dimensional fermion field is

$$L_{\text{int}} = g \psi(x_\mu) \psi(x_\mu) \Phi(x_\mu, x_4 = 0)$$

One can say that fermion field lives on 4-dimensional brane while scalar field lives in 5-dimensional world.

- For such interaction Feynman rules the standard as for 4-dimensional model except the use of effective scalar propagator

$$D^{\text{eff}}(k^2) = (2\pi)^{-1} \int [k^2 - f(k^2_4) + i\epsilon]^{-1} dk_4$$

One of possible generalizations to the gauge fields is to consider Yang-Mills in 4-dimensional space-time with standard action and matter fields in 5-dimensional space-time with the replacement of

the mass $m^2 \rightarrow f(-\partial_4^2)$. So for such kind of models gauge field $A_\mu^a(x)$ lives on four-dimensional brane, while matter field lives in 5-dimensional dimensional space-time and the Poincare invariance holds only in 4-dimensional space-time.

$$S_F = \int d^5x [\psi (i\gamma^\mu \partial_\mu + gT^a A^a_\mu \gamma^\mu - m(-\partial^2_4)) \psi]$$

Feynman rules for such model coincide with standard except the use of fermion propagator

$$i[\gamma^\mu p_\mu - m(p^2_4)]^{-1} \text{ and additional}$$

integration $(2\pi)^{-1} dp_4$ in fermion loop. For

the case when $m(p^2_4) = 0$ for $|p_4| < \varepsilon\pi$ and

$m(p^2_4) = \infty$ for $|p_4| > \varepsilon\pi$ the single difference

between our model and 4-dimensional case is

additional factor ε for each fermion loop due to additional integration over dp_4 in

fermion loop so the model is renormalizable and one loop β -function is

$$\beta(g) = -g^3(11N/3 - 2\varepsilon/3)/16\pi^2 + O(g^5)$$

Phenomenological implications

There are a lot of possible extensions of Standard Model with continuously distributed Higgs boson mass. For instance, consider SM in the unitary gauge and make replacement in free Higgs boson propagator

$$(p^2 - m_H^2)^{-1} \rightarrow D_{\text{int}}(p^2) = \int \rho(t) [p^2 - t + i\varepsilon]^{-1} dt$$

For $D_{\text{int}}(p^2) = (p^2 - m_H^2 + i\Gamma_{\text{int}} m_H)^{-1}$ we can interpret Γ_{int} as internal Higgs boson decay width into 5-th dimension.

For large $\Gamma_{\text{int}} \gg \Gamma_{\text{tot,H}}$ we shall have additional suppression factor

$$\Gamma_{\text{tot,H}} (\Gamma_{\text{tot,H}} + \Gamma_{\text{int}})^{-1}$$

for standard signatures like $pp \rightarrow H + \dots \rightarrow \gamma\gamma + \dots$

to be used at the LHC that can make the LHC Higgs boson discovery practically impossible.

Phenomenological implications

For $D_{\text{int}}(p^2) = \sum |c_n|^2 (p^2 - m_n^2 + i\varepsilon)^{-1}$ and
for $(m_k - m_{k-1})$ bigger than detector resolution we
shall have several peaks
in the reactions

$$pp \rightarrow \gamma\gamma + \dots$$

$$pp \rightarrow ZZ^* + \dots \rightarrow 4l + \dots$$

to be used for Higgs boson search at the
LHC with factor $|c_n|^2$ suppression for each
resonance that for big n makes the Higgs boson
discovery at LHC extremely difficult or even
impossible

Phenomenological implications

- It should be stressed that the proposed generalization of the SM model is renormalizable if the ultraviolet asymptotics of the Higgs boson propagator $D_{\text{int}}(p^2)$ coincides with free propagator

$$D_0(p^2) = (p^2)^{-1}$$

Phenomenological implications

Another possible implications are models of Z' bosons with continuously distributed mass. Most models predict the existence of new narrow vector boson Z' with

Total decay width $\Gamma_{\text{tot}} = O(10^{-2})M_{Z'}$, while in model with continuously distributed Z' boson mass Z' boson could be very broad and possible consequence is the existence of broad structure for dimuon mass distribution in the reaction



Conclusions

1. Unparticles can be interpreted as fields with continuously distributed mass.
2. Fields with continuously distributed mass can be treated as fields in $d > 4$ space-time and from experimental point of view it is not necessary to require Poincare group in D -dimensional space-time (only 4-dimensional Poincare group follow from experiment)
3. Renormalizable extensions at $d > 4$ are possible.
4. There are possible testable at the LHC phenomenological consequences like Higgs boson or Z' boson decaying into additional dimension(s)

